

OPEN

Five-month-old infants detect affiliation in colughter

Athena Vouloumanos¹ & Gregory A. Bryant²

Colughter—simultaneous laughter between two or more individuals—allows listeners across different cultures and languages to quickly evaluate affiliation within a social group. We examined whether infants are sensitive to acoustic information in colughter that indicates affiliation, specifically whether they can differentiate colughter between friends and colughter between strangers. In the first experiment, infants who heard alternating trials of colughter between friends and strangers listened longer to colughter between friends. In the second experiment, we examined whether infants were sensitive to the social context that was appropriate for each type of colughter. Infants heard colughter between friends and colughter between strangers preceded by a silent visual scene depicting one of two different social contexts: either two people affiliating or turning away from each other. Infants looked longer when the social scene was incongruent with the type of colughter. By 5 months, infants preferentially listen to colughter between friends and detect when colughter does not match the valence of a social interaction. The ability to rapidly evaluate acoustic features in colughter that reveal social relationships between novel individuals appears early in human infancy and might be the product of an adaptive affiliation detection system that uses vocal cues.

Infants begin engaging with the social environment and communicating with their caregivers within their first months of life. But infants also start observing the pattern of communicative and social behaviors between people around them^{1–4}. Through infants' third-party observations of social interactions, infants recognize affiliative intentions in social agents and use these to make predictions about others' behavior (for review see⁵).

Infants use a variety of types of information to reason about social relationships. For example, by 9 months, infants use shared positive evaluations, such as two people liking the same food, as a cue to affiliation^{6,7}. Infants at this age also infer that people who speak the same language will likely be friends⁸. Even younger infants can use an agent's helping or hindering behavior to infer the nature of the relationship—affiliative or antagonistic—between agents^{2,9}. In their second year, establishing affiliative relationships between agents through labels or actions creates expectations about how those agents will behave towards others^{10–12}, allowing infants to make sense of complex social relationships.

Our understanding of infants' inferences about others' social behaviors comes in part from how infants perceive visual cues of affiliative interaction, such as being hindered, harmed, or helped in some attempted action. But less is known about auditory cues of affiliative interaction. Vocal communicative behavior reveals a great deal about people's intentions, and infants are highly sensitive to intentional vocal signals such as speech e.g.,^{13,14}, but little research has examined how infants might use nonverbal auditory cues to make social inferences^{15,16}. Vocalizations are a particularly expressive communicative medium, including not only speech, the main channel for language use, but also emotional signaling¹⁵.

Laughter is a ubiquitous, nonverbal vocalization with a variety of functions, including communicating one's intentions to positively affiliate^{17,18}. Spontaneous laughter is generated by an evolutionarily conserved vocal emotion production system shared by most mammals^{19,20}. Human laughter is homologous to play vocalizations in nonhuman primates, and has been suggested to be a ritualized signal of heavy breathing during rough and tumble play¹⁷. Chimpanzees produce laugh-like play vocalizations to prolong playing time, suggesting that chimpanzees use laughter to signal they are engaging in non-threatening playing behavior that would otherwise resemble actual fighting²¹. Laughter in humans could serve a similar function as a non-threatening signal during tickling and rough and tumble play in children¹⁷.

But, in humans, volitional laughter has taken on a more complex and varied set of functions, such that it can manifest itself in almost any social context. In many social interactions, people often laugh together (i.e.,

¹Department of Psychology, New York University, New York, New York, USA. ²Department of Communication, University of California at Los Angeles, Los Angeles, California, USA. Correspondence and requests for materials should be addressed to A.V. (email: athena.vouloumanos@nyu.edu)




Figure 1. Waveform (top) and narrowband spectrogram (bottom; 30 ms Gaussian analysis window, 44.1 kHz sampling rate, 0–5 kHz frequency range) of friends' and strangers' laughter. Waveform representation shows speakers 1 and 2 on their respective channels. Blue lines represent fundamental frequency (F0) values (75–750 Hz range).

simultaneously), and this colughter varies with their friendship status²². When brief segments of colughter between friends and strangers were presented to listeners from 24 different societies, ranging from small-scale hunter-gatherers to rural farmers to urban students, listeners everywhere were able to distinguish colughter between friends from colughter between strangers²³. Acoustic analysis showed that listeners' judgments of affiliative status were largely driven by vocal features associated with physiological arousal²³. Specifically, laughter between friends was more likely to consist of shorter calls (i.e., faster sounding), less regular pitch and intensity cycles, and less variation in pitch cycle regularity (See Fig. 1). These acoustic differences are similar to the features that distinguish spontaneous and volitional laughter, and suggest that friends are more likely to engage in shared spontaneous laughter than strangers, something listeners can detect. Here we explore whether infants can use vocal information to infer affiliative status by distinguishing between laughter between friends and laughter between strangers. Spontaneous laughter has a long evolutionary history suggesting that humans may be particularly attuned early in development to the acoustic features associated with emotional vocal production.

Infants attend to others' laughter, and can use adults' laughter to understand others' actions and intentions. Six-month-olds use laughter for social referencing, for example looking at their parents more when they used an object unconventionally (using a ball as a clown nose) while laughing, than when they used it conventionally (bouncing a ball on the ground) while laughing²⁴. Infants who themselves laughed at the unconventional act looked at their parent less, suggesting they didn't need affective support. Infants also understand some of the functional significance of laughter, linking a person's vocal signals of humor to that person's intentions and actions. For instance, 15-month-old infants matched a person's humorous vocal signals (laughter and a humorous sentence) to her humorous actions¹⁶. Infants can thus recognize that laughter can be informative about a person's behavior.

But little work has examined how infants perceive laughter, and no work, to our knowledge, has explored whether infants can draw inferences about affiliative relationships from auditory information alone. In two experiments, we examined whether infants were sensitive to the acoustic and social signals in colughter between friends and strangers. In the first experiment, infants heard alternating trials of laughter between friends and laughter between strangers. We expected that infants would discriminate between the two types of colughter, and would attend preferentially to friends' colughter over strangers' colughter. In the second experiment, infants heard either colughter between friends or colughter between strangers, preceded by visual scenes depicting one of two different social contexts: either two actors affiliating with each other or turning away from each other (See Fig. 2). A positive affiliative condition was contrasted with a negative rather than a neutral condition to ensure that infants could differentiate and recognize the valence of the two interactions, consistent with prior studies examining infants' perceptions of social relationships^{6–8,25}. Also consistent with these prior infant studies on social relationships, we predicted that if infants recognize the social context that was appropriate to each type of laughter, they would look longer when the laughter and social context were incongruent: when stranger colughter followed affiliative interactions, and when friend colughter followed disengaged interactions.




Figure 2. Black and white still image from the affiliative video (A) and the disengaged video (B) shown in Experiment 2.

Experiment 1: Preference for Friend or Stranger Colaughter

Method. *Participants.* Participants were 24 healthy, full term infants (12 females; M age: 5 months 4 days, $SD = 12$ days) recruited from maternity wards at local hospitals (New York, NY). Parents reported the ethnicity of their infants as White (13), Black (3), Asian (1), mixed race (4), or chose not to answer (3). Of these infants, 5 were identified as Hispanic or Latino, 18 were identified as non-Hispanic and 1 family chose not to answer. Parents reported their highest educational degree as some college (3), college degree (4), graduate degree (16), or chose not to answer (1). An additional 10 infants were excluded because they looked at the screen for the maximum trial length for more than 50% of the trials (5), fussiness (2), a preexisting medical condition (1), or looking times greater than 2.5 standard deviations from the mean (2). Parents gave informed consent on behalf of their infants and received a certificate and small toys or t-shirts as gifts for their participation. All procedures were approved by the IRB at New York University (IRB-FY2016-81). All experiments were performed in accordance with the relevant guidelines and regulations.

Stimuli. Pretest and posttest music: Before and after the experimental trials, infants saw a black and white checkerboard while hearing a clip of Bach's Concerto for Violin and Orchestra No. 1 in A minor (BWV 1041-III. Allegro Assai). This music trial familiarized infants with the visual and sound aspects of the procedure e.g.,²⁶ so that we did not have to exclude any experimental data from the analyses e.g.,²⁷.

Colaughter segments were extracted from conversations between pairs of American English-speaking undergraduate students who participated in exchange for course credit. Participants either signed up with a friend whom they had known for any amount of time, or they signed up individually and were paired with a stranger. The participants were instructed to talk about any topic they chose. The average conversation length was 13.5 min (M length = 809.2 s, $SD = 151.3$). Each conversation participant wore a lapel microphone approximately 15 cm from their mouth (Sony ECM-77B) and was recorded digitally to DAT (16-bit amplitude resolution, 44.1-kHz sampling rate, uncompressed wav files, Sony DTC series recorder) on a separate audio channel. See²⁸ for a complete description of the conversations.

Based on previous work²³, we defined colaughter as concurrent laughter production (with intensity onsets within 1 s). The laughs could be either voiced (periodic) or unvoiced (aperiodic). Acoustically, laughter is variable but often characterized by an initial alerting component, stable vowel configurations, and decaying loudness and pitch^{29,17,30}. From 24 conversations, two colaughter segments were extracted from each (the first and last occurrence) for a total of 48 colaughter segments. Half of the conversations were between friends (M length of acquaintance = 20.5 months; Range = 4–54 months; M age = 18.6 years; $SD = 0.6$) and half were between newly acquainted strangers (M age = 19.3 years, $SD = 1.8$). An analysis of basic acoustic features was performed using Praat (version 5.4.15)³¹. F0 values were calculated using an autocorrelation method with recommended pitch

Acoustic measure	Colaughter type			
	Friends		Strangers	
Burst number	4.2	(1.8)	3.5	(1.6)
Bout length (ms)	1146	(455)	1067	(266)
Average burst duration (ms)	274	(108)	301	(99)
Laughter onset asynchrony	337	(299)	290	(209)
Mean F0 (Hz)	283	(88)	254	(79)
F0 SD (Hz)	43	(20)	32	(16)
F0 Min (Hz)	207	(67)	200	(65)
F0 Max (Hz)	377	(123)	329	(118)
F0 Range (Hz)	170	(88)	129	(77)
Center of gravity (Hz)	973	(342)	821	(427)
Loudness (dB)	59.6	(7.6)	60.3	(6.0)
Harmonics-to-noise ratio (dB)	5.5	(2.2)	6.4	(2.3)

Table 1. Acoustic analyses of friend colaughter and stranger colaughter. Values reported are means with standard deviations in parentheses. Laugh bursts were counted as a combination of the two speakers. For example, simultaneous laugh bursts between two speakers counted as one burst, but if two overlapping bursts were perceptible as two bursts, they were counted as two.

settings of 100–600 Hz for females and 75–500 Hz for males. Spectral measures were calculated using a cross correlation method. See Fig. 1 and Table 1 for acoustic characteristics. See²³ for more information about the colaughter segments.

Using these 48 colaughter segments (24 friend colaughter and 24 stranger colaughter), we created 4 audio files of friend colaughter and 4 audio files of stranger colaughter using Audacity 2.1.2 (Freeware distributed under GNU General Public License). Each trial included 12 laughter segments and was 24.5 s long with 300–1200 ms of silence between laughter segments.

Procedure. Infants were tested in a sound-attenuated room using an infant-controlled sequential preferential looking procedure (e.g.,^{27,26} run in Habit 2.1.25³²). In this procedure, infants controlled the onset and offset of each trial by looking at or away from a central monitor. Infants sat on a caregiver's lap 35'' (89 cm) in front of a 30'' (76.25 cm) computer monitor.

At the start of the experiment, infants' attention was drawn to the monitor by a colorful expanding and contracting circle. Once infants fixated on the monitor, a stationary black and white checkerboard appeared in tandem with one set of sounds, either friend colaughter or stranger colaughter, presented at a mean amplitude of 60 dB (± 5 dB). Sounds played until infants looked away from the monitor for 2 consecutive seconds, at which time the sounds and the visual display ceased.

The colorful expanding circle drew infants' attention back to the monitor between trials. Once infants fixated on the monitor, the stationary black and white checkerboard appeared in tandem with the other set of sounds. We presented 5 trials each of friend colaughter and stranger colaughter in alternation for a total of 10 experimental trials. Half the infants heard friend colaughter first and half the infants heard stranger colaughter first.

We compared infants' looking time to the screen during friend colaughter and stranger colaughter. Offline coders who were not aware of experimental condition did reliability coding on 20% of trials with the sound turned off using SuperCoder (Universal)³³ and reliability was high ($r = 0.99$).

Results. Infants listened longer to friend colaughter than stranger colaughter (see Fig. 3). A laughter (2: friend, stranger) by order (2: friend first, stranger first) by trial (5) by sex (2: female, male) ANOVA with age as a covariate revealed a significant effect of laughter, with infants listening longer to colaughter between friends ($M = 10.7$ s, $SD = 4.7$) than colaughter between strangers ($M = 9.6$, $SD = 4.1$), $F(1,19) = 4.93$, $p = 0.039$, partial $\eta^2 = 0.21$. A post-hoc power analysis showed that, based on this effect size of $f = 0.52$, and a sample of 24 infants, we had 66% power to detect a significant difference at an alpha level of $p < 0.05$ ³⁴. There were no other main effects or interactions.

Experiment 2: Detecting Social Context of Colaughter

Method. *Participants.* Participants were 24 healthy, full term infants (8 females; M age: 5 months 11 days, $SD = 11$ days) recruited from maternity wards at local hospitals (New York, NY). Parents reported the ethnicity of their infants as White (15), mixed race (6), or chose not to answer (3). Of these infants, 2 were identified as Hispanic or Latino, 14 identified as non-Hispanic and 8 families chose not to answer. Parents reported their highest educational degree as finished high school (1), college degree (8), graduate degree (11) or chose not to answer (4). An additional 19 infants were excluded because they looked at the screen for the maximum trial length for more than 50% of the trials (11), a preexisting medical condition (1), technical problems (2), experimenter error (4), or looking times greater than 2.5 standard deviations from the mean (1). This higher than expected attrition rate was driven by the very large number of infants (11) who did not disengage from the display on more than half of the trials and therefore could not give a reliable differential looking time measure. Parents gave informed consent on behalf of their infants and received a certificate and small toys or t-shirts as gifts for their participation. All procedures were approved by the IRB at New York University (IRB-FY2016-81).




Figure 3. Results. Individual (circles) and mean (blue line) difference scores of listening times (in seconds) to friend colaughter and stranger colaughter in Experiment 1. * indicates a significant difference at $p < 0.05$.

Stimuli. Visual displays consisted of 2-part videos with 2 female actors: In the first part, the actors engaged in a silent 6-s interaction in which they either acted affiliatively or disengaged towards each other (See Fig. 2). In the second part, which was presented immediately after the first part, the actors faced forward in a still frame (identical in both conditions) while co-laughter played. In the affiliative video sequence, the actors began side-by-side facing forward, turned toward each other, smiled and waved, and the scene faded to black. In the second part, the actors were shown in a still frame, facing forward toward the infant while either the congruent friend colaughter or the incongruent stranger colaughter played. In the disengaged video sequence, the actors began side-by-side facing forward, turned toward each other, immediately turned their back on each other and crossed their arms, and the scene faded to black. In the second part, the actors were shown in a still frame, facing forward toward the infant while either the congruent stranger colaughter or the incongruent friend colaughter played. The same auditory stimuli as in experiment 1 were used.

Procedure. Each trial had two parts: (1) a 6-s fixed length segment during which infants saw either the affiliative interaction or the disengaged interaction between the two actors, (2) a 24-s infant-controlled segment during which infants saw a still frame of the two actors facing forward while one of the laughter files was played. Infants either saw the affiliative or the disengaged interaction for each of the 10 experimental trials but the type of laughter alternated such that each infant heard 5 trials of friend colaughter and 5 trials of stranger colaughter. We compared infants' looking time to the screen during socially congruent trials (affiliative interaction followed by friend colaughter, disengaged interaction followed by stranger laughter) with socially incongruent trials (affiliative interaction followed by stranger colaughter, disengaged interaction followed by friend laughter). Offline coders who were not aware of experimental condition did reliability coding on 20% of trials with the sound turned off using SuperCoder (Universal)³³ and reliability was high ($r = 0.93$).

Results. Infants listened longer to incongruent trials than congruent trials, suggesting they noticed the discrepancy between cues in the social interaction and cues in the laughter vocalizations (see Fig. 4). A congruency (2: match, mismatch) by scene (2: affiliative, disengaged) by order (4: friend first, stranger first) by trial (5) by sex (2: female, male) ANOVA with age as a covariate revealed a significant effect of congruency, with infants listening longer to incongruent laughter ($M = 14.0$ s, $SD = 4.0$) than congruent laughter ($M = 12.6$, $SD = 4.8$), $F(1,19) = 4.92$, $p = 0.039$, partial $\eta^2 = 0.21$, and a significant effect of trial, with infants listening less in later trials, $F(4,16) = 5.28$, $p = 0.001$, partial $\eta^2 = 0.22$. There were no other main effects or interactions. A post-hoc power analysis showed that, as in Experiment 1, based on this effect size of $f = 0.52$, and a sample of 24 infants, we had 66% power to detect a significant difference at an alpha level of $p < 0.05$ ³⁴. There were no other main effects or interactions.

General Discussion

Vocal signals can quickly allow listeners to infer affiliative status between multiple individuals. Laughter constitutes an evolutionarily ancient, cross-culturally recognized social signal of affiliative status between people e.g.,²³. We found that 5-month-old infants prefer to listen to colaughter between friends over colaughter between strangers. A different group of infants recognized the social context that was appropriate to each type of laughter, and looked




Figure 4. Results. Individual (circles) and mean (blue line) difference scores of listening times (in seconds) to congruent and incongruent vocal and social cues in colughter in Experiment 2. * indicates a significant difference at $p < 0.05$.

longer when the type of laughter and social context were incongruent. These results provide the first evidence that infants can use a nonverbal vocal signal of affiliation in making social judgments as a third-party observer.

Infants' sensitivity to the social information in laughter by 5 months suggests early emerging, and possibly evolutionarily conserved, perceptual machinery for evaluating potential social signals that may be used to quickly draw inferences about the relationships between agents in the social environment. These findings are consistent with a growing body of work showing that infants attend to a range of information in establishing the social relationships between agents e.g.,^{11,12,6,8,2,35}.

Laughter is rich with acoustic features that infants could have used as the basis for their preference in Experiment 1. Both sets of colughter were used in a previous study²³, and an extensive acoustic analysis revealed that colughter judged as being between friends had more characteristics associated with speaker arousal, including frequency and temporal irregularities, and faster call rate. The descriptive acoustic analysis performed for the current study revealed that friends' colughter was higher in mean F_0 and F_0 variability, with also slightly higher maximum F_0 values (see Table 1). These F_0 properties (perceived as pitch) are also shared by infant-directed speech that adults often use to address infants, and that expresses positive emotion to infants and adults^{27,36,37}. Friend colughter may have thus sounded more positive in affect than stranger colughter to infants, and could provide a proximate explanation for why infants preferred friend colughter in Experiment 1. During the test trials of Experiment 2, infants heard laughter while seeing a still image of the actors, which suggests that those actors were not currently producing the laughter infants heard. This may suggest a more basic association between the visual scene and the laughter, rather than expecting the actors to laugh in a particular way. Moreover, infants might be sensitive to the positive affect common to both the affiliative scene and the colughter between friends, providing them with a basis for matching the social information in the two modalities in Experiment 2. This kind of perceptual sensitivity allows infants to begin to make predictions about important social information about others, such as who cooperates, and who is allied.

Whereas acoustic features in laughter can be informative about social affiliation, not all vocalizations appear to be informative about others' social relationships. For example, hearing others' conversation does not allow adults to predict their future actions in a behavioral economic game³⁸. Conversely, colughter potentially signals rich information about people's relationships, and is associated with actual cooperative behavior, at least in males³⁹, as well as being associated with other indicators of social engagement^{38,40}. Infants are sensitive to voices^{41,42}, and to particular tones of voices that signal positive affect and intentions to communicate (e.g., infant-directed speech and song⁴³⁻⁴⁵) and, like adults may be able to use acoustic features in speech to infer social relationships.

The current work reveals that infants can extract meaningful information from laughter about others' social relationships. Our findings suggest that infants can distinguish specific acoustic properties of laughter that index others' social relationships (as friends or strangers) and can match colughter to the affiliative status (affiliative or disengaged) of the interlocutors. Infants' sensitivity to different kinds of laughter might be one of the early emerging tools they use to understand and navigate the complex social world.

Data Availability

Data are publicly available on the Open Science Framework <https://osf.io/b43c8>.

References

1. Hamlin, J. K., Wynn, K., Bloom, P. & Mahajan, N. How infants and toddlers react to antisocial others. *Proc Natl Acad Sci USA* **108**, 19931–19936 (2011).
2. Hamlin, J. K., Wynn, K. & Bloom, P. Social evaluation by preverbal infants. *Nature* **450**, 557–559 (2007).
3. Vouloumanos, A., Martin, A. & Onishi, K. H. Do 6-month-olds understand that speech can communicate? *Dev Sci* **17**, 872–879 (2014).
4. Martin, A., Onishi, K. H. & Vouloumanos, A. Understanding the abstract role of speech in communication at 12 months. *Cognition* **123**, 50–60 (2012).
5. Van de Vondervoort, J. W. & Hamlin, J. K. The early emergence of sociomoral evaluation: infants prefer prosocial others. *Curr Opin Psychol* **20**, 77–81 (2018).
6. Liberman, Z., Kinzler, K. D. & Woodward, A. L. Friends or foes: infants use shared evaluations to infer others' social relationships. *J Exp Psychol Gen* **143**, 966–971 (2014).
7. Liberman, Z., Woodward, A. L., Sullivan, K. R. & Kinzler, K. D. Early emerging system for reasoning about the social nature of food. *Proc Natl Acad Sci USA* **113**, 9480–9485 (2016).
8. Liberman, Z., Woodward, A. L. & Kinzler, K. D. Preverbal infants infer third-party social relationships based on language. *Cogn Sci* **41**(Suppl 3), 622–634 (2017).
9. Hamlin, J. K., Wynn, K. & Bloom, P. Three-month-olds show a negativity bias in their social evaluations. *Dev Sci* **13**, 923–929 (2010).
10. Bian, L., Sloane, S. & Baillargeon, R. Infants expect ingroup support to override fairness when resources are limited. *Proc Natl Acad Sci USA* **115**, 2705–2710 (2018).
11. Jin, K. S. & Baillargeon, R. Infants possess an abstract expectation of ingroup support. *Proc Natl Acad Sci USA* **114**, 8199–8204 (2017).
12. Rhodes, M., Hetherington, C., Brink, K. & Wellman, H. M. Infants' use of social partnerships to predict behavior. *Dev Sci* **18**, 909–916 (2015).
13. Bryant, G. A. & Barrett, H. C. Recognizing intentions in infant-directed speech: evidence for universals. *Psychol Sci* **18**, 746–751 (2007).
14. Vouloumanos, A., Onishi, K. H. & Pogue, A. Twelve-month-old infants recognize that speech can communicate unobservable intentions. *Proc Natl Acad Sci USA* **109**, 12933–12937 (2012).
15. Sakkalou, E. & Gattis, M. Infants infer intentions from prosody. *Cogn Dev* **27**, 1–16 (2012).
16. Hoicka, E. & Wang, S.-H. Fifteen-month-old infants match vocal cues to intentional actions. *J Cogn Dev* **12**, 299–314 (2011).
17. Provine, R. R. *Laughter: A Scientific Investigation* (Viking, New York, 2000).
18. Bryant, G. A. & Aktipis, C. A. The animal nature of spontaneous human laughter. *Evol Hum Behav* **35**, 327–335 (2014).
19. Ackermann, H., Hage, S. R. & Ziegler, W. Brain mechanisms of acoustic communication in humans and nonhuman primates: an evolutionary perspective. *Behav Brain Sci* **37**, 529–546 (2014).
20. Jürgens, U. Neural pathways underlying vocal control. *Neurosci Biobehav Rev* **26**, 235–258 (2002).
21. Davila-Ross, M., Allcock, B., Thomas, C. & Bard, K. A. Aping expressions? Chimpanzees produce distinct laugh types when responding to laughter of others. *Emotion* **11**, 1013–1020 (2011).
22. Smoski, M. & Bachorowski, J. A. Antiphonal laughter between friends and strangers. *Cogn Emot* **17**, 327–340 (2003).
23. Bryant, G. A. *et al.* Detecting affiliation in colughter across 24 societies. *Proc Natl Acad Sci USA* **113**, 4682–4687 (2016).
24. Mireault, G. C. *et al.* Social looking, social referencing and humor perception in 6- and 12-month-old infants. *Infant Behav Dev* **37**, 536–545 (2014).
25. Liberman, Z., Kinzler, K. D. & Woodward, A. L. The early social significance of shared ritual actions. *Cognition* **171**, 42–51 (2018).
26. Vouloumanos, A. & Werker, J. F. Tuned to the signal: the privileged status of speech for young infants. *Dev Sci* **7**, 270–276 (2004).
27. Cooper, R. P. & Aslin, R. N. Developmental differences in infant attention to the spectral properties of infant-directed speech. *Child Dev* **65**, 1663–1677 (1994).
28. Bryant, G. A. Prosodic contrasts in ironic speech. *Discourse Process* **47**, 545–566 (2010).
29. Bachorowski, J.-A. & Owren, M. J. Not all laughs are alike: voiced but not unvoiced laughter readily elicits positive affect. *Psychol Sci* **12**, 252–257 (2001).
30. Titze, I. R., Finnegan, E. M., Laukkanen, A. M., Fujia, M. & Hoffman, H. Laryngeal muscle activity in giggle: a damped oscillation model. *J Voice* **22**, 644–648 (2008).
31. Boersma, P. & Weenink, D. Praat: doing phonetics by computer, Version 5.4.12, 2015. *Disponible en*. *Acesso em* **2** (2015).
32. Oakes, L. M., Sperka, D. J. & Cantrell, L. Habit 2. (Version 2.1.25). *Software* (2015).
33. Hollich, G. Supercoder: A program for coding preferential looking (Version 1.7.1). *Software* (2008).
34. Faul, F., Erdfelder, E., Lang, A. G. & Buchner, A. G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods* **39**, 175–191 (2007).
35. Hamlin, J. K. & Wynn, K. Young infants prefer prosocial to antisocial others. *Cogn Dev* **26**, 30–39 (2011).
36. Fernald, A. Intonation and communicative intent in mothers' speech to infants: is the melody the message? *Child Dev* **60**, 1497–1510 (1989).
37. Hoicka, E. Parents and toddlers distinguish joke, pretend and literal intentional contexts through communicative and referential cues. *J Pragmat* **95**, 137–155 (2016).
38. Manson, J. H., Gervais, M. M. & Kline, M. A. Defectors cannot be detected during "small talk" with strangers. *PLoS One* **8**, e82531 (2013).
39. Manson, J. H., Bryant, G. A., Gervais, M. M. & Kline, M. A. Convergence of speech rate in conversation predicts cooperation. *Evol Hum Behav* **34**, 419–426 (2013).
40. Dale, R., Bryant, G. A., Manson, J. H. & Gervais, M. M. Body synchrony in triadic conversation. (under review).
41. Grossmann, T., Oberecker, R., Koch, S. P. & Friederici, A. D. The developmental origins of voice processing in the human brain. *Neuron* **65**, 852–858 (2010).
42. Vouloumanos, A. & Werker, J. F. Listening to language at birth: evidence for a bias for speech in neonates. *Dev Sci* **10**, 159–164 (2007).
43. Fernald, A. In *The Adapted Mind: Evolutionary Psychology and the Generation of Culture* (eds Barkow, J. H. & Cosmides, L.) 391–428 (Oxford University Press, New York, NY, USA, 1992).
44. Paquette-Smith, M. & Johnson, E. K. I don't like the tone of your voice: Infants use vocal affect to socially evaluate others. *Infancy* **21**, 104–121 (2016).
45. Mehr, S. A., Song, L. A. & Spelke, E. S. For 5-Month-Old Infants, Melodies Are Social. *Psychol Sci* **27**, 486–501 (2016).

Acknowledgements

This research was funded by the Eunice Kennedy Shriver National Institute of Child Health & Human Development of the National Institutes of Health under Award Number R01HD072018 awarded to AV. We thank all of the members of the NYU Infant Cognition and Communication Lab and especially all of the families who participated in this study.

Author Contributions

A.V. and G.B. conceptualized the idea for the study. G.B. provided auditory stimuli and acoustic analyses and A.V. created visual stimuli. A.V. collected data and performed data analyses. A.V. and G.B. wrote the manuscript.

Additional Information

Competing Interests: The authors declare no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019